

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Fluzone® Quadrivalent safely and effectively. See full prescribing information for Fluzone Quadrivalent.

Fluzone Quadrivalent (Influenza Virus Vaccine)
Suspension for Intramuscular Injection
2013-2014 Formula
Initial US Approval (Fluzone Quadrivalent): 2013

INDICATIONS AND USAGE

Fluzone Quadrivalent is a vaccine indicated for active immunization for the prevention of influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine. (1)

Fluzone Quadrivalent is approved for use in persons 6 months of age and older. (1)

DOSAGE AND ADMINISTRATION

- For intramuscular use only (2)

Age	Dose	Schedule
6 months through 35 months	One or two doses ^a , 0.25 mL each	If 2 doses, administer at least 4 weeks apart
36 months through 8 years	One or two doses ^a , 0.5 mL each	If 2 doses, administer at least 4 weeks apart
9 years and older	One dose, 0.5 mL	-

^a1 or 2 doses depends on vaccination history as per Advisory Committee on Immunization Practices annual recommendations on prevention and control of influenza with vaccines

"-" Indicates information is not applicable

DOSAGE FORMS AND STRENGTHS

Suspension for injection supplied in 3 presentations: prefilled single-dose syringe (yellow plunger rod), 0.25 mL; prefilled single-dose syringe (purple plunger rod), 0.5 mL; single-dose vial, 0.5 mL. (3)

CONTRAINDICATIONS

Severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine, including egg protein, or after previous dose of any influenza vaccine. (4)

WARNINGS AND PRECAUTIONS

- If Guillain-Barré syndrome (GBS) has occurred within 6 weeks of previous influenza vaccination, the decision to give Fluzone Quadrivalent should be based on careful consideration of the potential benefits and risks. (5.1)

ADVERSE REACTIONS

- In children 6 months through 35 months of age, the most common ($\geq 10\%$) injection-site reactions were pain (57%) or tenderness (54%), erythema (37%), and swelling (22%); the most common solicited systemic adverse reactions were irritability (54%), abnormal crying (41%), malaise (38%), drowsiness (38%), appetite loss (32%), myalgia (27%), vomiting (15%), and fever (14%). (6.1)
- In children 3 years through 8 years of age, the most common ($\geq 10\%$) injection-site reactions were pain (67%), erythema (34%), and swelling (25%); the most common solicited systemic adverse reactions were myalgia (39%), malaise (32%), and headache (23%). (6.1)
- In adults 18 years and older, the most common ($\geq 10\%$) injection-site reaction was pain (47%); the most common solicited systemic adverse reactions were myalgia (24%), headache (16%), and malaise (11%). (6.1)
- In adults 65 years of age and older, the most common ($\geq 10\%$) injection-site reaction was pain (33%); the most common solicited systemic adverse reactions were myalgia (18%), headache (13%), and malaise (11%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sanofi Pasteur Inc., at 1-800-822-2463 (1-800-VACCINE) or VAERS at 1-800-822-7967 or www.vaers.hhs.gov.

USE IN SPECIFIC POPULATIONS

- Safety and effectiveness of Fluzone Quadrivalent have not been established in pregnant women or children less than 6 months of age. (8.1) (8.4)
- Pregnancy: Pregnancy registry available. Call Sanofi Pasteur Inc. at 1-800-822-2463. (8.1)
- Antibody responses to Fluzone Quadrivalent are lower in persons ≥ 65 years of age than in younger adults. (8.5)

See 17 FOR PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Issue date: June 2013

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
 - Dose and Schedule
 - Administration
- DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
 - Guillain-Barré Syndrome
 - Preventing and Managing Allergic Reactions
 - Altered Immunocompetence
 - Limitations of Vaccine Effectiveness
- ADVERSE REACTIONS
 - Clinical Trials Experience
 - Post-Marketing Experience
- USE IN SPECIFIC POPULATIONS
 - Pregnancy
 - Nursing Mothers
 - Pediatric Use
 - Geriatric Use

- DESCRIPTION
- CLINICAL PHARMACOLOGY
 - Mechanism of Action
- CLINICAL STUDIES
 - Immunogenicity of Fluzone Quadrivalent in Children 6 Months through 8 Years of Age
 - Immunogenicity of Fluzone Quadrivalent in Adults ≥ 18 years of age
 - Immunogenicity of Fluzone Quadrivalent in Geriatric Adults ≥ 65 years of age
- REFERENCES
- HOW SUPPLIED/STORAGE AND HANDLING
 - How Supplied
 - Storage and Handling
- PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

1 **FULL PRESCRIBING INFORMATION:**

2 **1 INDICATIONS AND USAGE**

3 Fluzone® Quadrivalent is an inactivated quadrivalent influenza virus vaccine indicated for the
4 prevention of influenza disease caused by influenza A subtype viruses and type B viruses
5 contained in the vaccine.

6

7 Fluzone Quadrivalent is approved for use in persons 6 months of age and older.

8

9 **2 DOSAGE AND ADMINISTRATION**

- 10 • For intramuscular use only

11 **2.1 Dose and Schedule**

12 The dose and schedule for Fluzone Quadrivalent are presented in [Table 1](#).

13 **Table 1: Dose and Schedule for Fluzone Quadrivalent**

Age	Dose	Schedule
6 months through 35 months	One or two doses ^a , 0.25 mL each	If 2 doses, administer at least 4 weeks apart
36 months through 8 years	One or two doses ^a , 0.5 mL each	If 2 doses, administer at least 4 weeks apart
9 years and older	One dose, 0.5 mL	-

14 ^a1 or 2 doses depends on vaccination history as per Advisory Committee on Immunization Practices annual
15 recommendations on prevention and control of influenza with vaccines

16 "-" Indicates information is not applicable

17

18 **2.2 Administration**

1 Inspect Fluzone Quadrivalent visually for particulate matter and/or discoloration prior to
2 administration. If any of these defects or conditions exist, the vaccine should not be administered.

3

4 Before administering a dose of vaccine, shake the prefilled syringe or single-dose vial. Withdraw
5 the vaccine using a sterile needle and syringe.

6

7 The preferred sites for intramuscular injection are the anterolateral aspect of the thigh in infants 6
8 months through 11 months of age, the anterolateral aspect of the thigh (or the deltoid muscle if
9 muscle mass is adequate) in persons 12 months through 35 months of age, or the deltoid muscle in
10 persons ≥ 36 months of age. The vaccine should not be injected into the gluteal area or areas
11 where there may be a major nerve trunk.

12

13 Do not administer this product intravenously, intradermally, or subcutaneously.

14

15 Fluzone Quadrivalent vaccine should not be combined through reconstitution or mixed with any
16 other vaccine.

17

18 **3 DOSAGE FORMS AND STRENGTHS**

19 Fluzone Quadrivalent is a suspension for injection.

20

21 Fluzone Quadrivalent is supplied in 3 presentations:

22 1) Prefilled single-dose syringe (yellow syringe plunger rod), 0.25 mL, for persons 6 months
23 through 35 months of age.

1 2) Prefilled single-dose syringe (purple syringe plunger rod), 0.5 mL, for persons 36 months of
2 age and older.

3 3) Single-dose vial, 0.5 mL, for persons 36 months of age and older.
4

5 **4 CONTRAINDICATIONS**

6 Do not administer Fluzone Quadrivalent to anyone with a history of a severe allergic reaction
7 (e.g., anaphylaxis) to any component of the vaccine [see [Description \(11\)](#)], including egg protein,
8 or to a previous dose of any influenza vaccine.
9

10 **5 WARNINGS AND PRECAUTIONS**

11 **5.1 Guillain-Barré Syndrome**

12 The 1976 swine influenza vaccine was associated with an elevated risk of Guillain-Barré
13 syndrome (GBS). Evidence for a causal relation of GBS with other influenza vaccines is
14 inconclusive; if an excess risk exists, it is probably slightly more than 1 additional case per 1
15 million persons vaccinated. (see ref. 1) If GBS has occurred within 6 weeks of previous influenza
16 vaccination, the decision to give Fluzone Quadrivalent should be based on careful consideration
17 of the potential benefits and risks.
18

19 **5.2 Preventing and Managing Allergic Reactions**

20 Appropriate medical treatment and supervision must be available to manage possible anaphylactic
21 reactions following administration of Fluzone Quadrivalent.
22

23 **5.3 Altered Immunocompetence**

1 If Fluzone Quadrivalent is administered to immunocompromised persons, including those
2 receiving immunosuppressive therapy, the expected immune response may not be obtained.

3

4 **5.4 Limitations of Vaccine Effectiveness**

5 Vaccination with Fluzone Quadrivalent may not protect all recipients.

6

7 **6 ADVERSE REACTIONS**

8 In children 6 months through 35 months of age, the most common ($\geq 10\%$) injection-site reactions
9 were pain (57%)^a or tenderness (54%)^b, erythema (37%), and swelling (22%); the most common
10 solicited systemic adverse reactions were irritability (54%), abnormal crying (41%), malaise
11 (38%), drowsiness (38%), appetite loss (32%), myalgia (27%), vomiting (15%), and fever
12 (14%). In children 3 years through 8 years of age, the most common ($\geq 10\%$) injection-site
13 reactions were pain (67%), erythema (34%), and swelling (25%); the most common solicited
14 systemic adverse reactions were myalgia (39%), malaise (32%), and headache (23%). In adults 18
15 years and older, the most common ($\geq 10\%$) injection-site reaction was pain (47%); the most
16 common solicited systemic adverse reactions were myalgia (24%), headache (16%), and malaise
17 (11%). In adults 65 years of age and older, the most common ($\geq 10\%$) injection-site reaction was
18 pain (33%); the most common solicited systemic adverse reactions were myalgia (18%), headache
19 (13%), and malaise (11%).

20

^a Assessed in children 24 months through 35 months of age

^b Assessed in children 6 months through 23 months of age

1 **6.1 Clinical Trials Experience**

2 Because clinical trials are conducted under widely varying conditions, adverse event rates
3 observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trial
4 of another vaccine, and may not reflect the rates observed in practice.

5

6 **Children 6 Months Through 8 Years of Age**

7 Study 1 (NCT01240746, see <http://clinicaltrials.gov>) was a single-blind, randomized, active-
8 controlled multi-center safety and immunogenicity study conducted in the US. In this study,
9 children 6 months through 35 months of age received one or two 0.25 mL doses of either Fluzone
10 Quadrivalent or one of two formulations of a comparator trivalent influenza vaccine (TIV-1 or
11 TIV-2), and children 3 years through 8 years of age received one or two 0.5 mL doses of either
12 Fluzone Quadrivalent, TIV-1, or TIV-2. Each of the trivalent formulations contained an influenza
13 type B virus that corresponded to one of the two type B viruses in Fluzone Quadrivalent (a type B
14 virus of the Victoria lineage or a type B virus of the Yamagata lineage). For participants who
15 received two doses, the doses were administered approximately 4 weeks apart. The safety analysis
16 set included 1841 children 6 months through 35 months of age and 2506 children 3 years through
17 8 years of age. Among participants 6 months through 8 years of age in the three vaccine groups
18 combined, 49.3% were female (Fluzone Quadrivalent, 49.2%; TIV-1, 49.8%; TIV-2, 49.4%),
19 58.4% Caucasian (Fluzone Quadrivalent, 58.4%; TIV-1, 58.9%; TIV-2, 57.8%), 20.2% Black
20 (Fluzone Quadrivalent, 20.5%; TIV-1, 19.9%; TIV-2, 19.1%), 14.1% Hispanic (Fluzone
21 Quadrivalent, 14.3%; TIV-1, 13.2%; TIV-2, 14.7%), and 7.3% were of other racial/ethnic groups
22 (Fluzone Quadrivalent, 6.8%; TIV-1, 8.0%; TIV-2, 8.5%). [Table 2](#) and [Table 3](#) summarize
23 solicited injection-site and systemic adverse reactions reported within 7 days post-vaccination via

1 diary cards. Participants were monitored for unsolicited adverse events for 28 days after each dose
2 and serious adverse events (SAEs) during the 6 months following the last dose.

3 **Table 2: Study 1^a: Percentage of Solicited Injection-site and Systemic Adverse Reactions**
4 **Within 7 Days After Vaccination in Children 6 Months Through 35 Months of Age (Safety**
5 **Analysis Set)^b**

	Fluzone Quadrivalent (N ^c =1223)			TIV-1 ^c (B Victoria) (N ^c =310)			TIV-2 ^d (B Yamagata) (N ^c =308)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
Injection-site adverse reactions									
- Pain ^h	57.0	10.2	1.0	52.3	11.5	0.8	50.3	5.4	2.7
- Tenderness ⁱ	54.1	11.3	1.9	48.4	8.2	1.9	49.7	10.3	0.0
- Erythema	37.3	1.5	0.2	32.9	1.0	0.0	33.3	1.0	0.0
- Swelling	21.6	0.8	0.2	19.7	1.0	0.0	17.3	0.0	0.0
Systemic adverse reactions									
- Fever (≥100.4° F) ^j	14.3	5.5	2.1	16.0	6.6	1.7	13.0	4.1	2.0
- Malaise ^h	38.1	14.5	4.6	35.2	14.8	4.7	32.4	12.8	6.8
- Myalgia ^h	26.7	6.6	1.9	26.6	9.4	1.6	25.0	6.8	2.7
- Headache ^h	8.9	2.5	0.6	9.4	3.9	0.0	12.2	4.7	0.0
- Irritability ⁱ	54.0	26.4	3.2	52.8	20.1	3.1	53.5	22.9	2.8
- Crying abnormal ⁱ	41.2	12.3	3.3	36.5	8.2	1.9	29.9	10.4	2.1
- Drowsiness ⁱ	37.7	8.4	1.3	32.1	3.8	0.6	31.9	5.6	0.7
- Appetite loss ⁱ	32.3	9.1	1.8	33.3	5.7	1.9	25.0	8.3	0.7
- Vomiting ⁱ	14.8	6.2	1.0	11.3	4.4	0.6	13.9	6.3	0.0

6 ^aNCT01240746

7 ^bThe safety analysis set includes all persons who received at least one dose of study vaccine

8 ^c2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
9 B/Brisbane/60/2008 (Victoria lineage), licensed

10 ^dInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
11 (Yamagata lineage), non-licensed

12 ^eN is the number of participants in the safety analysis set

- 1 ^fGrade 2 - Injection-site pain: sufficiently discomforting to interfere with normal behavior or activities; Injection-site
2 tenderness: cries and protests when injection-site is touched; Injection-site erythema, Injection-site swelling: ≥ 2.5
3 cm to < 5 cm; Fever: $> 101.3^{\circ}\text{F}$ to $\leq 103.1^{\circ}\text{F}$ (6 months through 23 months); $\geq 101.2^{\circ}\text{F}$ to $\leq 102.0^{\circ}\text{F}$ (24 months
4 through 35 months); Malaise, Myalgia, and Headache: some interference with activity; Irritability: requiring
5 increased attention; Crying abnormal: 1 to 3 hours; Drowsiness: not interested in surroundings or did not wake up
6 for a feed/meal; Appetite lost: missed 1 or 2 feeds/meals completely; Vomiting: 2 to 5 episodes per 24 hours
7
8 ^gGrade 3 - Injection-site pain: incapacitating, unable to perform usual activities; Injection-site tenderness: cries when
9 injected limb is moved, or the movement of the injected limb is reduced; Injection-site erythema, Injection-site
10 swelling: ≥ 5 cm; Fever: $> 103.1^{\circ}\text{F}$ (6 months through 23 months); $\geq 102.1^{\circ}\text{F}$ (24 months through 35 months);
11 Malaise, Myalgia, and Headache: Significant; prevents daily activity; Irritability: inconsolable; Crying abnormal: > 3
12 hours; Drowsiness: sleeping most of the time or difficult to wake up; Appetite lost: refuses ≥ 3 feeds/meals or
13 refuses most feeds/meals; Vomiting: ≥ 6 episodes per 24 hours or requiring parenteral hydration
14
15 ^hAssessed in children 24 months through 35 months of age
16 ⁱAssessed in children 6 months through 23 months of age
17 ^jFever measured by any route

17 **Table 3: Study 1^a: Percentage of Solicited Injection-site and Systemic Adverse Reactions**
18 **Within 7 Days After Vaccination in Children 3 Years Through 8 Years of Age (Safety**
19 **Analysis Set)^b**

	Fluzone Quadrivalent (N ^c =1669)			TIV-1 ^c (B Victoria) (N ^c =424)			TIV-2 ^d (B Yamagata) (N ^c =413)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
Injection-site adverse reactions									
- Pain	66.6	15.8	2.1	64.6	9.5	2.0	63.8	11.6	2.8
- Erythema	34.1	2.9	1.8	36.8	3.4	1.2	35.2	2.5	1.8
- Swelling	24.8	2.8	1.4	25.4	1.5	1.2	25.9	2.5	1.8
Systemic adverse reactions									
- Fever ($\geq 100.4^{\circ}\text{F}$) ^h	7.0	2.1	2.1	7.1	2.2	1.2	7.6	2.8	0.8
- Headache	23.1	6.8	2.2	21.2	5.1	2.7	24.4	7.5	2.0
- Malaise	31.9	11.2	5.5	32.8	11.4	5.6	33.4	10.8	5.0
- Myalgia	38.6	12.2	3.3	34.1	9.0	2.7	38.4	11.1	2.8

20 ^aNCT01240746

21 ^b The safety analysis set includes all persons who received at least one dose of study vaccine

22 ^c2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
23 B/Brisbane/60/2008 (Victoria lineage), licensed

^dInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006 (Yamagata lineage), non-licensed

^eN is the number of participants in the safety analysis set

^fGrade 2 - Injection-site pain: sufficiently discomforting to interfere with normal behavior or activities; Injection-site erythema, Injection-site swelling: ≥ 2.5 cm to < 5 cm; Fever: $\geq 101.2^{\circ}\text{F}$ to $\leq 102.0^{\circ}\text{F}$; Headache, Malaise, and Myalgia: some interference with activity

^gGrade 3 - Injection-site pain: incapacitating, unable to perform usual activities; Injection-site erythema, Injection-site swelling: ≥ 5 cm; Fever: $\geq 102.1^{\circ}\text{F}$; Headache, Malaise, and Myalgia: Significant; prevents daily activity

^hFever measured by any route

Among children 6 months through 8 years of age, unsolicited non-serious adverse events were reported in 1360 (47.0%) recipients in the Fluzone Quadrivalent group, 352 (48.0%) recipients in the TIV-1 group, and 346 (48.0%) recipients in the TIV-2 group. The most commonly reported unsolicited non-serious adverse events were cough, vomiting, and pyrexia. During the 28 days following vaccination, a total of 16 (0.6%) recipients in the Fluzone Quadrivalent group, 4 (0.5%) recipients in the TIV-1 group, and 4 (0.6%) recipients in the TIV-2 group, experienced at least one SAE; no deaths occurred. Throughout the study period, a total of 41 (1.4%) recipients in the Fluzone Quadrivalent group, 7 (1.0%) recipients in the TIV-1 group, and 14 (1.9%) recipients in the TIV-2 group, experienced at least one SAE. Three SAEs were considered to be possibly related to vaccination: croup in a Fluzone Quadrivalent recipient and 2 episodes of febrile seizure, 1 each in a TIV-1 recipient and a TIV-2 recipient. One death occurred in the TIV-1 group (a drowning 43 days post-vaccination).

Adults

In study 2 (NCT00988143, see <http://clinicaltrials.gov>), a multi-centered randomized, open-label trial conducted in the US, adults 18 years of age and older received one dose of either Fluzone Quadrivalent or one of two formulations of comparator trivalent influenza vaccine (TIV-1 or TIV-2). Each of the trivalent formulations contained an influenza type B virus that corresponded to one

1 of the two type B viruses in Fluzone Quadrivalent (a type B virus of the Victoria lineage or a type
2 B virus of the Yamagata lineage). The safety analysis set included 570 recipients, half aged 18-60
3 years and half aged 61 years or older. Among participants in the three vaccine groups combined,
4 67.2% were female (Fluzone Quadrivalent, 68.4%; TIV-1, 67.9%; TIV-2, 65.3%), 88.4%
5 Caucasian (Fluzone Quadrivalent, 91.1%; TIV-1, 86.8%; TIV-2, 87.4%), 9.6% Black (Fluzone
6 Quadrivalent, 6.8%; TIV-1, 12.1%; TIV-2, 10.0%), 0.4% Hispanic (Fluzone Quadrivalent, 0.0%;
7 TIV-1, 0.5%; TIV-2, 0.5%), and 1.7% were of other racial/ethnic groups (Fluzone Quadrivalent,
8 2.1%; TIV-1, 0.5%; TIV-2, 2.2%). [Table 4](#) summarizes solicited injection-site and systemic
9 adverse reactions reported within 3 days post-vaccination via diary cards. Participants were
10 monitored for unsolicited adverse events and SAEs during the 21 days following vaccination.

11 **Table 4: Study 2^a: Percentage of Solicited Injection-site and Systemic Adverse Reactions**
12 **Within 3 Days After Vaccination in Adults 18 Years of Age and Older (Safety Analysis Set)^b**

	Fluzone Quadrivalent (N ^c =190)			TIV-1 ^c (B Victoria) (N ^c =190)			TIV-2 ^d (B Yamagata) (N ^c =190)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
Injection-site adverse reactions									
- Pain	47.4	6.8	0.5	52.1	7.9	0.5	43.2	6.3	0.0
- Erythema	1.1	0.0	0.0	1.6	0.5	0.0	1.6	0.5	0.0
- Swelling	0.5	0.0	0.0	3.2	0.5	0.0	1.1	0.0	0.0
- Induration	0.5	0.0	0.0	1.6	0.5	0.0	0.5	0.0	0.0
- Ecchymosis	0.5	0.0	0.0	0.5	0.0	0.0	0.5	0.0	0.0
Systemic adverse reactions									
- Myalgia	23.7	5.8	0.0	25.3	5.8	0.0	16.8	5.8	0.0
- Headache	15.8	3.2	0.5	18.4	6.3	0.5	18.0	4.2	0.0
- Malaise	10.5	1.6	1.1	14.7	3.2	1.1	12.1	4.7	0.5

	Fluzone Quadrivalent (N ^e =190)			TIV-1 ^c (B Victoria) (N ^e =190)			TIV-2 ^d (B Yamagata) (N ^e =190)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
- Shivering	2.6	0.5	0.0	5.3	1.1	0.0	3.2	0.5	0.0
- Fever (≥100.4° F) ^h	0.0	0.0	0.0	0.5	0.5	0.0	0.5	0.5	0.0

1 ^aNCT00988143

2 ^bThe safety analysis set includes all persons who received study vaccine

3 ^c2009-2010 Fluzone TIV containing A/Brisbane/59/2007 (H1N1), A/Uruguay/716/2007 (H3N2), and
4 B/Brisbane/60/2008 (Victoria lineage), licensed

5 ^d2008-2009 Fluzone TIV containing A/Brisbane/59/2007 (H1N1), A/Uruguay/716/2007 (H3N2), and
6 B/Florida/04/2006 (Yamagata lineage), licensed

7 ^eN is the number of participants in the safety analysis set

8 ^fGrade 2 - Injection-site pain: Some interference with activity; Injection-site erythema, Injection-site swelling,
9 Injection-site induration, and Injection-site ecchymosis: ≥5.1 to ≤10 cm; Fever: ≥101.2°F to ≤102.0°F; Myalgia,
10 Headache, Malaise, and Shivering: some interference with activity

11 ^gGrade 3 - Injection-site pain: Significant; prevents daily activity; Injection-site erythema, Injection-site swelling,
12 Injection-site induration, and Injection-site ecchymosis: >10 cm; Fever: ≥102.1°F; Myalgia, Headache, Malaise, and
13 Shivering: Significant; prevents daily activity

14 ^hFever measured by any route

15

16 Unsolicited non-serious adverse events were reported in 33 (17.4%) recipients in the Fluzone

17 Quadrivalent group, 45 (23.7%) recipients in the TIV-1 group, and 45 (23.7%) recipients in the

18 TIV-2 group. The most commonly reported unsolicited non-serious adverse events were

19 headache, cough, and oropharyngeal pain. In the follow-up period, there were two SAEs, 1 (0.5%)

20 in the Fluzone Quadrivalent group and 1 (0.5%) in the TIV-2 group. No deaths were reported

21 during the trial period.

22

23 Geriatric Adults

24 In Study 3 (NCT01218646, see <http://clinicaltrials.gov>), a multi-center, randomized, double-blind

25 trial conducted in the US, adults 65 years of age and older received one dose of either Fluzone

26 Quadrivalent, or one of two formulations of comparator trivalent influenza vaccine (TIV-1 or

1 TIV-2). Each of the trivalent formulations contained an influenza type B virus that corresponded
 2 to one of the two type B viruses in Fluzone Quadrivalent (a type B virus of the Victoria lineage or
 3 a type B virus of the Yamagata lineage). The safety analysis set included 675 recipients. Among
 4 participants in the three vaccine groups combined, 55.7% were female (Fluzone Quadrivalent,
 5 57.3%; TIV-1, 56.0%; TIV-2, 53.8%), 89.5% Caucasian (Fluzone Quadrivalent, 87.6%; TIV-1,
 6 89.8%; TIV-1, 91.1%), 2.2% Black (Fluzone Quadrivalent, 4.0%; TIV-1, 1.8%; TIV, 0.9%), 7.4%
 7 Hispanic (Fluzone Quadrivalent, 8.4%; TIV-1, 7.6%; TIV, 6.2%) and 0.9% were of other
 8 racial/ethnic groups (Fluzone Quadrivalent, 0.0%; TIV-1, 0.9%; TIV, 1.8%).

9
 10 [Table 5](#) summarizes solicited injection-site and systemic adverse reactions reported within 7 days
 11 post-vaccination via diary cards. Participants were monitored for unsolicited adverse events and
 12 SAEs during the 21 days following vaccination.

13
 14 **Table 5: Study 3^a: Percentage of Solicited Injection-site and Systemic Adverse Reactions**
 15 **Within 7 Days After Vaccination in Adults 65 Years of Age and Older (Safety Analysis Set)^b**

	Fluzone Quadrivalent (N ^c =225)			TIV-1 ^c (B Victoria) (N ^c =225)			TIV-2 ^d (B Yamagata) (N ^c =225)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
Injection-site adverse reactions									
- Pain	32.6	1.3	0.9	28.6	2.7	0.0	23.1	0.9	0.0
- Erythema	2.7	0.9	0.0	1.3	0.0	0.0	1.3	0.4	0.0
- Swelling	1.8	0.4	0.0	1.3	0.0	0.0	0.0	0.0	0.0
Systemic adverse reactions									
- Myalgia	18.3	4.0	0.4	18.3	4.0	0.0	14.2	2.7	0.4

	Fluzone Quadrivalent (N ^c =225)			TIV-1 ^c (B Victoria) (N ^c =225)			TIV-2 ^d (B Yamagata) (N ^c =225)		
	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)	Any (%)	Grade 2 ^f (%)	Grade 3 ^g (%)
- Headache	13.4	1.3	0.4	11.6	1.3	0.0	11.6	1.8	0.4
- Malaise	10.7	4.5	0.4	6.3	0.4	0.0	11.6	2.7	0.9
- Fever (≥100.4° F) ^h	1.3	0.0	0.4	0.0	0.0	0.0	0.9	0.4	0.4

1 ^aNCT01218646

2 ^b The safety analysis set includes all persons who received study vaccine

3 ^c2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
4 B/Brisbane/60/2008 (Victoria lineage), licensed

5 ^dInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
6 (Yamagata lineage), non-licensed

7 ^eN is the number of participants in the safety analysis set

8 ^fGrade 2 - Injection-site pain: some interference with activity; Injection-site erythema and Injection-site swelling:
9 ≥5.1 to ≤10 cm; Fever: ≥101.2°F to ≤102.0°F; Myalgia, Headache, and Malaise: some interference with activity

10 ^gGrade 3 - Injection-site pain: Significant; prevents daily activity ; Injection-site erythema and Injection-site swelling:
11 >10 cm; Fever: ≥102.1°F; Myalgia, Headache, and Malaise: Significant; prevents daily activity

12 ^hFever measured by any route

13

14 Unsolicited non-serious adverse events were reported in 28 (12.4%) recipients in the Fluzone
15 Quadrivalent group, 22 (9.8%) recipients in the TIV-1 group, and 22 (9.8%) recipients in the TIV-
16 2 group. The most commonly reported adverse events were oropharyngeal pain, rhinorrhea,
17 injection-site induration, and headache. Three SAEs were reported during the follow-up period, 2
18 (0.9%) in the TIV-1 group and 1 (0.4%) in the TIV-2 group. No deaths were reported during the
19 trial period.

20

21 **6.2 Post-Marketing Experience**

22 Currently, there are no post-marketing data available for Fluzone Quadrivalent vaccine.

23

1 The following events have been spontaneously reported during the post-approval use of the
2 trivalent formulation of Fluzone. Because these events are reported voluntarily from a population
3 of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal
4 relationship to vaccine exposure. Adverse events were included based on one or more of the
5 following factors: severity, frequency of reporting, or strength of evidence for a causal
6 relationship to Fluzone.

7

- 8 • *Blood and Lymphatic System Disorders*: Thrombocytopenia, lymphadenopathy
- 9 • *Immune System Disorders*: Anaphylaxis, other allergic/hypersensitivity reactions (including
10 urticaria, angioedema)
- 11 • *Eye disorders*: Ocular hyperemia
- 12 • *Nervous System Disorders*: Guillain-Barré syndrome (GBS), convulsions, febrile
13 convulsions, myelitis (including encephalomyelitis and transverse myelitis), facial palsy
14 (Bell's palsy), optic neuritis/neuropathy, brachial neuritis, syncope (shortly after vaccination),
15 dizziness, paresthesia
- 16 • *Vascular Disorders*: Vasculitis, vasodilatation/flushing
- 17 • *Respiratory, Thoracic and Mediastinal Disorders*: Dyspnea, pharyngitis, rhinitis, cough,
18 wheezing, throat tightness
- 19 • *Skin and Subcutaneous Tissue Disorders*: Stevens-Johnson syndrome
- 20 • *General Disorders and Administration Site Conditions*: Pruritus, asthenia/fatigue, pain in
21 extremities, chest pain
- 22 • *Gastrointestinal Disorders*: Vomiting

23

1 **8 USE IN SPECIFIC POPULATIONS**

2 **8.1 Pregnancy**

3 Pregnancy Category C: Animal reproduction studies have not been conducted with Fluzone
4 Quadrivalent. It is also not known whether Fluzone Quadrivalent can cause fetal harm when
5 administered to a pregnant woman or can affect reproduction capacity. Fluzone Quadrivalent
6 should be given to a pregnant woman only if clearly needed.

7

8 Sanofi Pasteur Inc. is conducting a prospective pregnancy exposure registry to collect data on
9 pregnancy outcomes and newborn health status following vaccination with Fluzone Quadrivalent
10 during pregnancy. Healthcare providers are encouraged to enroll women who receive Fluzone
11 Quadrivalent during pregnancy in Sanofi Pasteur Inc.'s vaccination pregnancy registry by calling
12 1-800-822-2463.

13

14 **8.3 Nursing Mothers**

15 It is not known whether Fluzone Quadrivalent is excreted in human milk. Because many drugs are
16 excreted in human milk, caution should be exercised when Fluzone Quadrivalent is administered
17 to a nursing woman.

18

19 **8.4 Pediatric Use**

20 Safety and effectiveness of Fluzone Quadrivalent in children below the age of 6 months have not
21 been established. Safety and immunogenicity of Fluzone Quadrivalent was evaluated in children 6
22 months through 8 years of age. [See *Adverse Reactions (6.1) and Clinical Studies (14.1)*]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

8.5 Geriatric Use

Safety and immunogenicity of Fluzone Quadrivalent was evaluated in adults 65 years of age and older. [See *Clinical Studies (14.3)*] Antibody responses to Fluzone Quadrivalent are lower in persons ≥ 65 years of age than in younger adults.

11 DESCRIPTION

Fluzone Quadrivalent (Influenza Virus Vaccine) for intramuscular injection is an inactivated influenza virus vaccine, prepared from influenza viruses propagated in embryonated chicken eggs.

The virus-containing allantoic fluid is harvested and inactivated with formaldehyde. Influenza virus is concentrated and purified in a linear sucrose density gradient solution using a continuous flow centrifuge. The virus is then chemically disrupted using a non-ionic surfactant, Octylphenol Ethoxylate (Triton® X-100), producing a “split virus”. The split virus is further purified and then suspended in sodium phosphate-buffered isotonic sodium chloride solution. The Fluzone Quadrivalent process uses an additional concentration factor after the ultrafiltration step in order to obtain a higher hemagglutinin (HA) antigen concentration. Antigens from the four strains included in the vaccine are produced separately and then combined to make the quadrivalent formulation.

Fluzone Quadrivalent suspension for injection is clear and slightly opalescent in color.

Neither antibiotics nor preservative are used in the manufacture of Fluzone Quadrivalent.

1 The Fluzone Quadrivalent prefilled syringe or vial presentations are not made with natural rubber
2 latex.
3
4 Fluzone Quadrivalent is standardized according to United States Public Health Service
5 requirements and is formulated to contain 60 micrograms (mcg) HA per 0.5 mL dose in the
6 recommended ratio of 15 mcg HA of each of the following four influenza strains recommended
7 for the 2013-2014 influenza season: A/California/07/2009 X-179A (H1N1), A/Texas/50/2012 X-
8 223A (H3N2) (an A/Victoria/361/2011-like virus), B/Massachusetts/02/2012 (B Yamagata
9 lineage), and B/Brisbane/60/2008 (B Victoria lineage). The amounts of HA and other ingredients
10 per dose of vaccine are listed in [Table 6](#).

1 **Table 6: Fluzone Quadrivalent Ingredients**

Ingredient	Quantity (per dose)	
	Fluzone Quadrivalent 0.25 mL Dose	Fluzone Quadrivalent 0.5 mL Dose
Active Substance: Split influenza virus, inactivated strains^a:	30 mcg HA total	60 mcg HA total
A (H1N1)	7.5 mcg HA	15 mcg HA
A (H3N2)	7.5 mcg HA	15 mcg HA
B/(Victoria lineage)	7.5 mcg HA	15 mcg HA
B/(Yamagata lineage)	7.5 mcg HA	15 mcg HA
Other:		
Sodium phosphate-buffered isotonic sodium chloride solution	QS ^b to appropriate volume	QS ^b to appropriate volume
Formaldehyde	≤50 mcg	≤100 mcg
Octylphenol Ethoxylate	≤125 mcg	≤250 mcg
Preservative	None	None

2 ^aper United States Public Health Service (USPHS) requirement

3 ^bQuantity Sufficient

4

5 **12 CLINICAL PHARMACOLOGY**

6 **12.1 Mechanism of Action**

7 Influenza illness and its complications follow infection with influenza viruses. Global surveillance
8 of influenza identifies yearly antigenic variants. Since 1977, antigenic variants of influenza A
9 (H1N1 and H3N2) viruses and influenza B viruses have been in global circulation. Since 2001,
10 two distinct lineages of influenza B (Victoria and Yamagata lineages) have co-circulated
11 worldwide. Protection from influenza virus infection has not been correlated with a specific level
12 of hemagglutination inhibition (HI) antibody titer post-vaccination. However, in some human
13 studies, antibody titers $\geq 1:40$ have been associated with protection from influenza illness in up to
14 50% of subjects. (see ref. 2) (see ref. 3)

1

2 Antibodies against one influenza virus type or subtype confer limited or no protection against
3 another. Furthermore, antibodies to one antigenic variant of influenza virus might not protect
4 against a new antigenic variant of the same type or subtype. Frequent development of antigenic
5 variants through antigenic drift is the virologic basis for seasonal epidemics and the reason for the
6 usual change of one or more new strains in each year's influenza vaccine. Therefore, influenza
7 vaccines are standardized to contain the hemagglutinins of influenza virus strains, representing the
8 influenza viruses likely to be circulating in the US in the upcoming winter.

9

10 Annual vaccination with the current vaccine is recommended because immunity during the year
11 after vaccination declines, and because circulating strains of influenza virus change from year to
12 year.

13

14 **14 CLINICAL STUDIES**

15 **14.1 Immunogenicity of Fluzone Quadrivalent in Children 6 Months through 8** 16 **Years of Age**

17 In Study 1 (NCT01240746) (see [Section 6.1](#)), 1419 children 6 months through 35 months of age
18 and 2101 children 3 years through 8 years of age were included in the per-protocol
19 immunogenicity analysis. Participants received one or two 0.25 mL doses or one or two 0.5 mL
20 doses, respectively of Fluzone Quadrivalent, TIV-1, or TIV-2. For participants who received two
21 doses, the doses were administered approximately 4 weeks apart. The distribution of demographic
22 characteristics was similar to that of the safety analysis (see [Section 6.1](#)).

23

1 HI antibody geometric mean titers (GMTs) and seroconversion rates 28 days following
2 vaccination with Fluzone Quadrivalent were non-inferior to those following each TIV for all four
3 strains, based on pre-specified criteria (see [Table 7](#) and [Table 8](#)).

4 **Table 7: Study 1^a: Non-inferiority of Fluzone Quadrivalent Relative to TIV for Each Strain**
5 **by HI Antibody GMTs at 28 Days Post-Vaccination, Persons 6 Months Through 8 Years of**
6 **Age (Per-protocol Analysis Set)^b**

Antigen Strain	Fluzone Quadrivalent N ^c =2339	Pooled TIV ^d N ^c =1181		GMT Ratio (95%CI) ^e
	GMT	GMT		
A (H1N1)	1124	1096		1.03 (0.93; 1.14)
A (H3N2)	822	828		0.99 (0.91; 1.08)
	Fluzone Quadrivalent N ^c =2339	TIV-1 ^f (B Victoria) N ^c =582	TIV-2 ^g (B Yamagata) N ^c =599	GMT Ratio (95%CI) ^e
	GMT	GMT	GMT	
B/Brisbane/60/2008 (B Victoria)	86.1	64.3	(19.5) ^h	1.34 (1.20; 1.50)
B/Florida/04/2006 (B Yamagata)	61.5	(16.3) ⁱ	58.3	1.06 (0.94; 1.18)

7 ^aNCT01240746

8 ^bPer-protocol analysis set included all persons who had no study protocol deviations

9 ^cN is the number of participants in the per-protocol analysis set

10 ^dPooled TIV group includes participants vaccinated with either TIV-1 or TIV-2

11 ^eNon-inferiority was demonstrated if the lower limit of the 2-sided 95% CI of the ratio of GMTs (Fluzone
12 Quadrivalent divided by pooled TIV for the A strains, or the TIV containing the corresponding B strain) was > 0.66

13 ^f2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
14 B/Brisbane/60/2008 (Victoria lineage), licensed

15 ^gInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
16 (Yamagata lineage), non-licensed

17 ^hTIV-2 did not contain B/Brisbane/60/2008

18 ⁱTIV-1 did not contain B/Florida/04/2006

19

1 **Table 8: Study 1^a: Non-inferiority of Fluzone Quadrivalent Relative to TIV for Each Strain**
 2 **by Seroconversion Rates at 28 Days Post-Vaccination, Persons 6 Months Through 8 Years**
 3 **of Age (Per-protocol Analysis Set)^b**

Antigen Strain	Fluzone Quadrivalent N ^c =2339	Pooled TIV ^d N ^c =1181		Difference of Seroconversion Rates (95% CI) ^f
	Seroconversion ^e (%)			
A (H1N1)	92.4	91.4		0.9 (-0.9; 3.0)
A (H3N2)	88.0	84.2		3.8 (1.4; 6.3)
	Fluzone Quadrivalent N ^c =2339	TIV-1 ^g (B Victoria) N ^c =582	TIV-2 ^h (B Yamagata) N ^c =599	Difference of Seroconversion Rates (95% CI) ^f
	Seroconversion ^e (%)			
B/Brisbane/60/2008 (B Victoria)	71.8	61.1	(20.0) ⁱ	10.7 (6.4; 15.1)
B/Florida/04/2006 (B Yamagata)	66.1	(17.9) ^j	64.0	2.0 (-2.2; 6.4)

4 ^aNCT01240746

5 ^bPer-protocol analysis set included all persons who had no study protocol deviations

6 ^cN is the number of participants in the per-protocol analysis set

7 ^dPooled TIV group includes participants vaccinated with either TIV-1 or TIV-2

8 ^eSeroconversion: Paired samples with pre-vaccination HI titer <1:10 and post-vaccination titer ≥1:40 or a minimum
 9 4-fold increase for participants with pre-vaccination titer ≥1:10

10 ^fNon-inferiority was demonstrated if the lower limit of the 2-sided 95% CI of the difference in seroconversion rates
 11 (Fluzone Quadrivalent minus pooled TIV for the A strains, or the TIV containing the corresponding B strain) was
 12 >-10%

13 ^g2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
 14 B/Brisbane/60/2008 (Victoria lineage), licensed

15 ^hInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
 16 (Yamagata lineage), non-licensed

17 ⁱTIV-2 did not contain B/Brisbane/60/2008

18 ^jTIV-1 did not contain B/Florida/04/2006

19

20 Non-inferiority immunogenicity criteria based on HI antibody GMTs and seroconversion rates

21 were also met when age subgroups (6 months to <36 months and 3 years to <9 years) were

22 examined. In addition, HI antibody GMTs and seroconversion rates following Fluzone

23 Quadrivalent were higher than those following TIV for the B strain not contained in each

24 respective TIV based on pre-specified criteria (the lower limit of the 2-sided 95% CI of the ratio

1 of the GMTs [Fluzone Quadrivalent divided by TIV] >1.5 for each B strain in Fluzone
2 Quadrivalent compared with the corresponding B strain not contained in each TIV and the lower
3 limit of the two 2-sided 95% CI of the difference of the seroconversion rates [Fluzone
4 Quadrivalent minus TIV] >10% for each B strain in Fluzone Quadrivalent compared with the
5 corresponding B strain not contained in each TIV).

6

7 **14.2 Immunogenicity of Fluzone Quadrivalent in Adults ≥ 18 Years of Age**

8 In Study 2 (NCT00988143) (see [Section 6.1](#)), 565 adults 18 years of age and older who had
9 received one dose of Fluzone Quadrivalent, TIV-1, or TIV-2 were included in the per-protocol
10 immunogenicity analysis. The distribution of demographic characteristics was similar to that of
11 the safety analysis (see [Section 6.1](#)).

12

13 HI antibody GMTs 21 days following vaccination with Fluzone Quadrivalent were non-inferior to
14 those following each TIV for all four strains, based on pre-specified criteria (see [Table 9](#)).

15 **Table 9: Study 2^a: Non-inferiority of Fluzone Quadrivalent Relative to TIV for Each Strain**
16 **by HI Antibody GMTs at 21 Days Post-Vaccination, Adults 18 Years of Age and Older (Per-**
17 **protocol Analysis Set)^b**

Antigen Strain	Fluzone Quadrivalent N ^c =190	Pooled TIV ^d N ^c =375		GMT Ratio (95%CI) ^e
	GMT	GMT		
A (H1N1)	161	151		1.06 (0.87, 1.31)
A (H3N2)	304	339		0.90 (0.70, 1.15)
	Fluzone Quadrivalent N ^c =190	TIV-1 ^f (B Victoria) N ^c =187	TIV-2 ^g (B Yamagata) N ^c =188	GMT Ratio (95%CI) ^e
	GMT	GMT	GMT	

Antigen Strain	Fluzone Quadrivalent N ^c =190	Pooled TIV ^d N ^c =375		GMT Ratio (95%CI) ^e
B/Brisbane/60/2008 (B Victoria)	101	114	(44.0) ^h	0.89 (0.70, 1.12)
B/Florida/04/2006 (B Yamagata)	155	(78.1) ⁱ	135	1.15 (0.93, 1.42)

1 ^aNCT00988143

2 ^bPer-protocol analysis set included all persons who had no study protocol deviations

3 ^cN: number of participants in the per-protocol analysis set

4 ^dPooled TIV group includes participants vaccinated with either TIV-1 or TIV-2

5 ^eNon-inferiority was demonstrated if the lower limit of the 2-sided 95% CI of the ratio of GMTs (Fluzone
6 Quadrivalent divided by pooled TIV for the A strains, or the TIV containing the corresponding B strain) was >2/3

7 ^f2009-2010 Fluzone TIV containing A/Brisbane/59/2007 (H1N1), A/Uruguay/716/2007 (H3N2), and
8 B/Brisbane/60/2008 (Victoria lineage), licensed

9 ^g2008-2009 Fluzone TIV containing A/Brisbane/59/2007 (H1N1), A/Uruguay/716/2007 (H3N2), and
10 B/Florida/04/2006 (Yamagata lineage), licensed

11 ^hTIV-2 did not contain B/Brisbane/60/2008

12 ⁱTIV-1 did not contain B/Florida/04/2006

13

14 **14.3 Immunogenicity of Fluzone Quadrivalent in Geriatric Adults ≥ 65 Years of** 15 **Age**

16 In Study 3 (NCT01218646) (see [Section 6.1](#)), 660 adults 65 years of age and older were included
17 in the per-protocol immunogenicity analysis set. The distribution of demographic characteristics
18 was similar to that of the safety analysis (see [Section 6.1](#)).

19

20 HI antibody GMTs 21 days following vaccination with Fluzone Quadrivalent were non-inferior to
21 those following TIV for all four strains, based on pre-specified criteria (see [Table 10](#)).

22 Seroconversion rates 21 days following Fluzone Quadrivalent were non-inferior to those
23 following TIV for H3N2, B/Brisbane, and B/Florida, but not for H1N1 (see [Table 11](#)). The HI
24 antibody GMT following Fluzone Quadrivalent was higher than that following TIV-1 for
25 B/Florida but not higher than that following TIV-2 for B/Brisbane, based on pre-specified criteria

1 (the lower limit of the 2-sided 95% CI of the ratio of the GMTs [Fluzone Quadrivalent divided by
2 TIV] >1.5 for each B strain in Fluzone Quadrivalent compared with the corresponding B strain
3 not contained in each TIV). Seroconversion rates following Fluzone Quadrivalent were higher
4 than those following TIV for the B strain not contained in each respective TIV, based on pre-
5 specified criteria (the lower limit of the two 2-sided 95% CI of the difference of the
6 seroconversion rates [Fluzone Quadrivalent minus TIV] >10% for each B strain in Fluzone
7 Quadrivalent compared with the corresponding B strain not contained in each TIV).

8 **Table 10: Study 3^a: Non-inferiority of Fluzone Quadrivalent Relative to TIV for Each Strain**
9 **by HI Antibody GMTs at 21 Days Post-Vaccination, Adults 65 Years of Age and Older (Per-**
10 **protocol Analysis Set)^b**

Antigen Strain	Fluzone Quadrivalent N ^c =220	Pooled TIV ^d N ^c =440		GMT Ratio (95%CI) ^e
	GMT	GMT		
A (H1N1)	231	270		0.85 (0.67; 1.09)
A (H3N2)	501	324		1.55 (1.25; 1.92)
	Fluzone Quadrivalent N ^c =220	TIV-1 ^f (B Victoria) N ^c =219	TIV-2 ^g (B Yamagata) N ^c =221	GMT Ratio (95%CI) ^e
	GMT	GMT	GMT	
B/Brisbane/60/2008 (B Victoria)	73.8	57.9	(42.2) ^h	1.27 (1.05; 1.55)
B/Florida/04/2006 (B Yamagata)	61.1	(28.5) ⁱ	54.8	1.11 (0.90; 1.37)

11 ^aNCT01218646

12 ^bPer-protocol analysis set included all persons who had no study protocol deviations

13 ^cN: number of participants in the per-protocol analysis set

14 ^dPooled TIV group includes participants vaccinated with either TIV-1 or TIV-2

15 ^eNon-inferiority was demonstrated if the lower limit of the 2-sided 95% CI of the ratio of GMTs (Fluzone
16 Quadrivalent divided by pooled TIV for the A strains, or the TIV containing the corresponding B strain) was >0.66

17 ^f2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
18 B/Brisbane/60/2008 (Victoria lineage), licensed

19 ^gInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
20 (Yamagata lineage), non-licensed

1 ^hTIV-2 did not contain B/Brisbane/60/2008
2 ⁱTIV-1 did not contain B/Florida/04/2006
3

4 **Table 11: Study 3^a: Non-inferiority of Fluzone Quadrivalent Relative to TIV for Each Strain**
5 **by Seroconversion Rates at 21 Days Post-Vaccination, Adults 65 Years of Age and Older**
6 **(Per-protocol Analysis Set)^b**

Antigen Strain	Fluzone Quadrivalent N ^c =220	Pooled TIV ^d N ^c =440		Difference of Seroconversion Rate (95% CI) ^e
	Seroconversion ^f (%)			
A (H1N1)	65.91	69.77		-3.86 (-11.50; 3.56)
A (H3N2)	69.09	59.32		9.77 (1.96; 17.20)
	Fluzone Quadrivalent N ^c =220	TIV-1 ^g (B Victoria) N ^c =219	TIV-2 ^h (B Yamagata) N ^c =221	Difference of Seroconversion Rate (95% CI) ^e
	Seroconversion ^f (%)			
B/Brisbane/60/2008 (B Victoria)	28.64	18.72	(8.60) ⁱ	9.91 (1.96; 17.70)
B/Florida/04/2006 (B Yamagata)	33.18	(9.13) ^j	31.22	1.96 (-6.73; 10.60)

7 ^aNCT01218646
8 ^bPer-protocol analysis set included all persons who had no study protocol deviations
9 ^cN: number of participants in the per-protocol analysis set
10 ^dPooled TIV group includes participants vaccinated with either TIV-1 or TIV-2
11 ^eNon-inferiority was demonstrated if the lower limit of the 2-sided 95% CI of the difference in seroconversion rates
12 (Fluzone Quadrivalent minus pooled TIV for the A strains, or the TIV containing the corresponding B strain) was
13 >-10%
14 ^fSeroconversion: Paired samples with pre-vaccination HI titer <1:10 and post-vaccination titer ≥1:40 or a minimum 4-
15 fold increase for participants with pre-vaccination titer ≥1:10
16 ^g2010-2011 Fluzone TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and
17 B/Brisbane/60/2008 (Victoria lineage), licensed
18 ^hInvestigational TIV containing A/California/07/2009 (H1N1), A/Victoria/210/2009 (H3N2), and B/Florida/04/2006
19 (Yamagata lineage), non-licensed
20 ⁱTIV-2 did not contain B/Brisbane/60/2008
21 ^jTIV-1 did not contain B/Florida/04/2006
22

1 **15 REFERENCES**

2

3 1 Lasky T, Terracciano GJ, Magder L, et al. The Guillain-Barré syndrome and the 1992-1993
4 and 1993-1994 influenza vaccines. *N Engl J Med* 1998;339:1797-802.

5 2 Hannoun C, Megas F, Piercy J. Immunogenicity and protective efficacy of influenza
6 vaccination. *Virus Res* 2004;103:133-138.

7 3 Hobson D, Curry RL, Beare AS, Ward-Gardner A. The role of serum haemagglutination-
8 inhibiting antibody in protection against challenge infection with influenza A2 and B
9 viruses. *J Hyg Camb* 1972;70:767-777.

10

1 **16 HOW SUPPLIED/STORAGE AND HANDLING**

2 **16.1 How Supplied**

3 Single-dose, prefilled syringe (yellow plunger rod), without needle, 0.25 mL (NDC 49281-513-
4 00) (not made with natural rubber latex). Supplied as package of 10 (NDC 49281-513-25).

5

6 Single-dose, prefilled syringe (purple plunger rod), without needle, 0.5 mL (NDC 49281-413-88)
7 (not made with natural rubber latex). Supplied as package of 10 (NDC 49281-413-50).

8

9 Single-dose vial, 0.5 mL (NDC 49281-413-58) (not made with natural rubber contain latex).

10 Supplied as package of 10 (NDC 49281-413-10).

11

12 **16.2 Storage and Handling**

13 Store all Fluzone Quadrivalent presentations refrigerated at 2° to 8°C (35° to 46°F). DO NOT
14 FREEZE. Discard if vaccine has been frozen.

15

16 Do not use after the expiration date shown on the label.

17

18 **17 PATIENT COUNSELING INFORMATION**

19 See FDA-approved patient labeling (Patient Information). Inform the vaccine recipient or

20 guardian:

- 21 • Fluzone Quadrivalent contains killed viruses and cannot cause influenza.
- 22 • Fluzone Quadrivalent stimulates the immune system to protect against influenza, but does not
23 prevent other respiratory infections.

- 1 • Annual influenza vaccination is recommended.
- 2 • Report adverse reactions to their healthcare provider and/or to the Vaccine Adverse Event
3 Reporting System (VAERS) at 1-800-822-7967.
- 4 • Sanofi Pasteur Inc. is conducting a prospective pregnancy exposure registry to collect data on
5 pregnancy outcomes and newborn health status following vaccination with Fluzone
6 Quadrivalent during pregnancy. Women who receive Fluzone Quadrivalent during pregnancy
7 are encouraged to contact Sanofi Pasteur Inc. directly or have their healthcare provider contact
8 Sanofi Pasteur Inc. at 1-800-822-2463.

9

10 Vaccine Information Statements must be provided to vaccine recipients or their guardians, as
11 required by the National Childhood Vaccine Injury Act of 1986 prior to immunization. These
12 materials are available free of charge at the Centers for Disease Control and Prevention (CDC)
13 website (www.cdc.gov/vaccines).

14

15 Fluzone is a registered trademark of Sanofi Pasteur Inc.

16

17 Manufactured by:

18 **Sanofi Pasteur Inc.**

19 Swiftwater PA 18370 USA

20

21



1 **Patient Information Sheet**
2 **Fluzone® Quadrivalent**
3 **Influenza Virus Vaccine**
4

5 Please read this information sheet before getting Fluzone® Quadrivalent vaccine. This summary is
6 not intended to take the place of talking with your healthcare provider. If you have questions or
7 would like more information, please talk with your healthcare provider.

8
9 **What is Fluzone Quadrivalent vaccine?**

10 Fluzone Quadrivalent is a vaccine that helps protect against influenza illness (flu).

11 Fluzone Quadrivalent vaccine is for people who are 6 months of age and older.

12 Vaccination with Fluzone Quadrivalent vaccine may not protect all people who receive the
13 vaccine.

14
15 **Who should not get Fluzone Quadrivalent vaccine?**

16 You should not get Fluzone Quadrivalent vaccine if you:

- 17 • ever had a severe allergic reaction to eggs or egg products.
18 • ever had a severe allergic reaction after getting any flu vaccine.
19 • are younger than 6 months of age.

20
21 Tell your healthcare provider if you or your child have or have had:

- 22 • Guillain-Barré syndrome (severe muscle weakness) after getting a flu vaccine.
23 • problems with your immune system as the immune response may be diminished.

24
25 **How is the Fluzone Quadrivalent vaccine given?**

26 Fluzone Quadrivalent vaccine is a shot given into the muscle of the arm.

1 For infants, Fluzone Quadrivalent vaccine is a shot given into the muscle of the thigh.

2

3 **What are the possible side effects of Fluzone Quadrivalent vaccine?**

4 The most common side effects of Fluzone Quadrivalent vaccine are:

- 5 • pain, redness, swelling, bruising and hardness where you got the shot
- 6 • muscle aches
- 7 • tiredness
- 8 • headache
- 9 • fever

10 These are not all of the possible side effects of Fluzone Quadrivalent vaccine. You can ask your
11 healthcare provider for a list of other side effects that is available to healthcare professionals.

12 Call your healthcare provider for advice about any side effects that concern you. You may report
13 side effects to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967 or
14 <http://vaers.hhs.gov>. Sanofi Pasteur Inc. is collecting information on pregnancy outcomes and the
15 health of newborns following vaccination with Fluzone Quadrivalent during pregnancy. Women
16 who receive Fluzone Quadrivalent during pregnancy are encouraged to contact Sanofi Pasteur Inc.
17 directly or have their healthcare provider contact Sanofi Pasteur Inc. at 1-800-822-2463.

18

19 **What are the ingredients in Fluzone Quadrivalent vaccine?**

20 Fluzone Quadrivalent vaccine contains 4 killed flu virus strains.

21 Inactive ingredients include formaldehyde and octylphenol ethoxylate.

22

23 Manufactured by:

1 **Sanofi Pasteur Inc.**
2 Swiftwater, PA 18370 USA
3

